

EXPLICIT DIAGONALIZATION OF THE MARKOV FORM ON THE TEMPERLEY-LIEB ALGEBRA

JOSH GENAUER AND NEAL W. STOLTZFUS

ABSTRACT. In fundamental paper in 1984, Vaughn Jones developed his new polynomial invariant of knots using a Markov trace on the Temperley-Lieb algebra. Subsequently, Lickorish used the associated bilinear pairing to provided an alternative proof for the existence of the 3-manifold invariants of Witten, Reshetinkin, and Turaev. A key property of this form is the non-degeneracy of this form except at the parameter values $\pm 2\cos\frac{\pi}{n+1}$ [7]. Ko and Smolinsky derived a recursive formula for the determinants of specific minors of Markov's form, establishing the needed non-degeneracy [6]. In this paper, we define a triangular change of basis in which the form is diagonal and explicitly compute the diagonal entries of this matrix as products of quotients of Chebyshev polynomials, corroborating the determinant computation of Ko and Smolinsky. The method of proof employs a recursive method for defining the required orthogonal basis elements in the Temperley-Lieb algebra, similar in spirit to Jones' and Wenzl's recursive formula for a family of projectors in the Temperley-Lieb algebra. We define a partial order on the non-crossing chord diagram basis and give an explicit formula for a recursive construction of an orthogonal basis, via a recursion over this partial order. Finally we relate this orthogonal basis to bases constructed using the calculus of trivalent graphs developed by Kauffman and Lins[5].

1. INTRODUCTION

In 1984, Vaughn Jones developed his new polynomial invariant of knots using a Markov trace, t , on the Temperley-Lieb algebra. The Markov trace can be extended to a bilinear pairing using the algebra structure, $\langle x, y \rangle = t(xy)$. In the basis given by non-crossing chord diagrams (disjoint arcs in the upper half plane with their endpoints on $\{1, 2, 3, \dots, 2n\}$), the pairing can be defined geometrically by joining the first non-crossing chord diagram with the reflection of the second non-crossing chord diagram in the lower half-plane, counting the number of circles, n , and associating to this pair of non-crossing chord diagrams, the monomial, q^n , where q is an indeterminate, called the *quantum parameter*. Using the theory of quantum groups and representations of lie algebras, Witten, Reshetinkin, and Turaev proved the existence of new invariants for three manifolds and links in three manifolds [8, 10]. Using the Markov associated bilinear form, Lickorish [7] gave an alternative proof of the existence of these quantum invariants of a 3-manifold.

A key property of this form is its non-degeneracy except when specialized at the real numbers, $\pm 2\cos\frac{\pi}{n+1}$, the real parts of complex roots of unity and the roots of the Chebychev polynomials [7]. Ko and Smolinsky derived a recursive formula for

Research partially supported by NSF Grant DMS-0353722, and a Louisiana Board of Regents Enhancement grant, LEQSF(2002-2004)-ENH-TR-17.

the determinants of Markov's form in the non-crossing chord diagram basis and established the non-degeneracy needed by Lickorish to complete his definition[6].

In this paper, we define a triangular change of basis to an orthogonal basis for the form and explicitly compute the diagonal entries of this matrix as products of quotients of Chebyshev polynomials. To accomplish this, we define a new partial order on the set of *non-crossing* chord diagrams (which form a geometric basis for the Temperley-Lieb algebra) and use this partial order to compute an orthogonal basis, by a recursion over this poset.

In a related setting of Topological Quantum Field Theory, Blanchet, Habegger, Masbaum and Vogel have exploited a related bilinear form and explicit orthogonal bases, see Lemma 4.1 in [1] Pat Gilmer has also communicated the fact that other orthogonal bases for the Temperley-Lieb algebra were previously known, see the comment in [3], and can be constructed using the method of trivalent graphs discussed in Kauffman and Lins[5]. In the penultimate section, we relate our orthogonal basis to those constructed by trivalent graph techniques.

2. PRELIMINARIES

Definition 2.1. A *non-crossing chord diagram* is an isotopy class of a set of n non-intersecting properly embedded arcs (a configuration) in the closed upper half-plane joining $2n$ points on the real line, the boundary of the upper half plane.

In this paper, we consider only non-crossing chord diagrams, as they provide a geometric basis for the Temperley-Lieb algebra. Henceforth, we drop the adjective, non-crossing. Note also that the isotopy, in the definition above, is not required to be fixed on the boundary, although it must be proper, that is, taking the boundary into itself. In each isotopy class, there is a unique chord diagram in *standard position* defined by the condition that the arcs are semicircles with centers on the real axis whose endpoints are located at the positive integers, $\{1, 2, 3, \dots, 2n\}$. We will often use this configuration for a chord diagram implicitly within our presentation. We adopt the perspective and notation of Ko and Smolinsky [6] for the set, D_n , of chord diagrams.

Remark 2.1. The cardinality of the set, D_n of n -chord diagrams is the n th Catalan number, $C_n = \frac{1}{n+1} \binom{2n}{n}$ [2].

There are many equivalent combinatorial reformulations for chord diagrams (see R. Stanley [9], for 63 variants of the ‘‘Catalan disease’’), but we will mention only two relevant equivalent descriptions. Graphically, we can uniquely associate to each chord diagram the *rooted plane tree* which is dual to the arcs (and rooted to the top). A second algebraic perspective is to associate the pairing of the end-points of a given non-intersecting chord diagram. This is a fixed-point-free involution (*perfect matching* in the combinatorial vernacular) on the set $\{1, 2, 3, \dots, 2n\}$ satisfying the condition that the orbit transpositions are non-crossing.

3. THE MARKOV SYMMETRIC BILINEAR FORM

Consider the ring $R = \mathbb{Z}[q, q^{-1}]$ and M , the free R -module generated by the chord diagrams. Let $\mathbb{Q}(q)$ be the quotient field of R , the field rational functions (quotients of polynomials with rational coefficients), and denote by V_n the free vector space spanned by D_n , that is, the set of formal finite sums of n -chord diagrams with coefficients in $\mathbb{Q}(q)$. A basis for this vector space will be denoted $e_{(a_n, \dots, a_1)}$

and ordered by the lexicographic order. Following Lickorish [7], we define a bilinear symmetric form: $\langle \cdot, \cdot \rangle_n : V_n \times V_n \rightarrow \mathbb{Q}(q)$ to be the bilinear extension of the geometric pairing on the chord diagram basis given by associating the monomial, q^c , where c is the count of the number of closed curves obtained when two chord diagrams are *glued* together.

Explicitly, we consider two n -chord diagram α and β in standard position with endpoints at $\{1, \dots, 2n\}$ on the real line $y = 0$ in the plane. Reflect β across the real axis, and then identify each node of α with its corresponding node in β (the correspondence is determined by connecting endpoints with the same index). This defines a configuration we will denote $\alpha \cup \beta$. Count the number of resulting circles in $\alpha \cup \beta$, and call this number $c := \#(\alpha \cup \beta)$. We then define $\langle \alpha, \beta \rangle_n := q^c$. This defines the Markov form on a linearly independent basis for V_n and we extend the form by bilinearity to all of $V_n \times V_n$. The following figure illustrates the construction:




FIGURE 1. $\langle (1, 3, 2, 1, 1), (2, 1, 2, 1, 1) \rangle = x^2$

4. CHEBYSHEV POLYNOMIALS

Let T_k denote the $k \times k$ tri-diagonal matrix:

$$\begin{pmatrix} q & 1 & & & \\ 1 & q & 1 & & 0 \\ & 1 & q & 1 & \\ & & \ddots & \ddots & \ddots \\ 0 & & & 1 & q & 1 \\ & & & & 1 & q \end{pmatrix}$$


and let the k th Chebyshev polynomial Δ_k be defined as $\det(T_k)$, a polynomial in q . By expanding the determinant along its top row we find the following recursive relation for $k \geq 3$:

$$(4.1) \quad \Delta_k = q\Delta_{k-1} - \Delta_{k-2}$$

Defining $\Delta_0 := 1$ and $\Delta_{-1} := 0$ the above identity holds for $k \geq 1$.

5. OPERATIONS ON n -CHORD DIAGRAMS

We also need the following operations on chord diagrams from Ko and Smolinsky. For $0 < k < 2n + 1$, let $l_k : D_n \rightarrow D_{n+1}$ be defined by taking a chord diagram in standard position and adding an arc in the upper half plane whose endpoints lie on the real line between the points $(k - 1, 0)$ and $(k, 0)$ so that the inserted arc is sufficiently small to be a new inner loop, i.e. it does not intersect any other arcs. Finally, we take the isotopy class of this new chord diagrams with $2n + 2$ endpoints in D_{n+1} .

FIGURE 2. l_5

For $0 < k < 2n + 1$, we also define: $\tau_k : D_{n+1} \rightarrow D_n$ by taking a chord diagram and adding the interval $[k, k + 1]$ on the x -axis. If k and $k + 1$ were already joined by an arc we remove the resulting component. The result is a chord diagram on $2n - 2$ vertices and we take its isotopy class in D_{n-1} .

Remark 5.1. Note that $\tau_j l_k$ is the identity on D_n for $j \in \{k - 1, k, k + 1\}$.



Remark 5.2. The number of arcs is finite, so that every chord diagram must have an innermost arc.

Finally, we extend the operations, l_k and τ_k linearly to the vector space, V , spanned by the chord diagrams.

Hence any $\alpha \in D_n$ there is a $\beta \in D_{n-1}$ such that $\alpha = l_k(\beta)$ for some k . Namely set $\beta = \tau_k(\alpha)$ where α has an innermost arc with endpoints k and $k + 1$. Moreover k is uniquely determined if we insist that k is as small as possible. Recursively we obtain a unique sequence $(l_{k_n}, l_{k_{n-1}}, \dots, l_{k_1})$ such that $\alpha = l_{k_n}(l_{k_{n-1}}(\dots(l_{k_1}(\phi))))$, where ϕ denotes the empty chord diagram with no arcs. For brevity's sake we will often omit the l 's from the n -tuple and refer only to the unique sequence $(k_n, \dots, k_1)$. Following Ko and Smolinsky, we call this tuple the *restricted* sequence associated to the chord diagram α .

Remark 5.3. Given an arbitrary sequence $(k_n, \dots, k_1)$ with $0 < k < 2n - 1$ one can associate the element of D_n given by $l_{k_n}(l_{k_{n-1}}(\dots(l_{k_1}(\phi))))$. This less restricted association is, however, not one-to-one, so we shall only use restricted sequences (unless explicitly noted otherwise).

The following simple characterization of restricted sequences will be quite useful.

FIGURE 3. τ_7 FIGURE 4. The chord diagram with associated restricted sequence $(3, 2, 2, 1, 2, 2, 1)$

Proposition 5.1. A sequence of positive integers $(k_n, \dots, k_1)$ is restricted if and only if $k_1 = 1$, $k_{i+1} \leq k_i + 1$ for $1 \leq i \leq n - 1$.

First note that, given a sequence $(k_n, \dots, k_1)$, the leftmost inner arc has its left endpoint at $(k_n, 0)$. The only constraint when adding a new arc is to guarantee that it is the leftmost inner arc, that is, its left endpoint must be in the range from $(1, 0)$ to $(k_n + 1, 0)$.

pen

6. COORDINATE-WISE PARTIAL ORDER ON RESTRICTED SEQUENCE

Using the restricted sequences, we will now define a coordinate-wise partial order on the chord diagrams.

Definition 6.1. $\alpha = (a_n, \dots, a_1) \preceq \beta = (b_n, \dots, b_1)$ is provided that, for all k satisfying $1 \leq k \leq n$, we have $a_k \leq b_k$.

The cover diagrams for these finite ranked posets are drawn below for the cases, $n = 3, 4$ and 5 .

Finally, we prove a proposition concerning this partial order that will be needed later.




FIGURE 5. Cover Diagram for the Poset of Non-crossing 3-chords




FIGURE 6. Cover Diagram for the Poset of Non-crossing 4-chords




FIGURE 7. Cover Diagram for the Poset of Non-crossing 5-chords

Proposition 6.1. Let $\beta \in D_n$ be a chord diagram with associated restricted sequence $(b_n, \dots, b_1)$. If $j \notin \{b_n - 1, b_n, b_n + 1\}$, then the restricted sequence associated to $\tau_j(\beta)$ is $(b'_{n-1}, \dots, b'_1)$ where:

- (1) $b'_{n-1} = b_n - 2$ if $j < b_n - 1$
- (2) $b'_{n-1} \leq b_n$ if $j > b_n + 1$.

Proof. For the first statement, note that the arcs whose endpoints are j and $j + 1$ must have the endpoints p and q to the right of the leftmost inner arc. Otherwise, there would be an inner arc to the left of the leftmost inner arc. Applying τ_j , the new chord diagram is obtained from the old by deleting two arcs and joining p and q by an arc which does not affect the leftmost inner arc's position. By deleting two arcs to the left of the inner arc, the inner arc has shifted two spots to the left. Thus $b'_{n-1} = b_n - 2$.

To verify second statement, we consider the endpoints of the arcs whose endpoints are at j and $j + 1$. If there is an inner loop at j (from j to $j + 1$), then removing that inner arc does nothing to affect the position of the leftmost inner arc. Then $b'_{n-1} = b_n$. Similarly if one arc or both arcs have their end points to the right of the leftmost inner arc, then τ_j does not change the position of the leftmost

inner arc. If both arcs have endpoints to the left of the leftmost inner arc, then these points must be j' and $j' + 1$ where $j' + 1 < b_n$ (they must be consecutive points, or else there would be an inner arc to the left of the leftmost inner arc). But now τ_j changes the chord diagram by deleting the two arcs at j and $j + 1$ and adding an arc from j' to $j' + 1$ and leaving the rest of the diagram unchanged. At the conclusion there is an inner arc to the left of b_n , so $b'_{n-1} < b_n$. $\square$

7. OPERATIONS & THE MARKOV FORM

We now state explicitly several basic facts relating the previously defined operations and the Markov form used implicitly in [6].

Proposition 7.1. If $\alpha, \beta \in D_n$, then for $1 \leq k \leq 2n + 1$, $\langle l_k(\alpha), l_k(\beta) \rangle = q \cdot \langle \alpha, \beta \rangle$.

Proof. Geometrically, the inserted arcs in $l_k(\alpha)$ and $l_k(\beta)$ join to form an additional circle disjoint from those formed by $\alpha \cup \beta$. $\square$




FIGURE 8. $\langle l_k(\alpha), l_k(\beta) \rangle$

Proposition 7.2. If $\alpha, \beta \in D_n$, then for $1 \leq k \leq 2n$, $\langle l_{k+1}(\alpha), l_k(\beta) \rangle = \langle \alpha, \beta \rangle$

Proof. Again, draw the picture. In this case the inserted arcs form an interval which is in some component of $\alpha \cup \beta$. The number of components is unchanged.




FIGURE 9. $\langle l_k(\alpha), l_{k+1}(\beta) \rangle$

$\square$

Upon linear extension to V , Proposition 7.1 and Proposition 7.2 imply:

Proposition 7.3. For $f, g \in V_n$ and $1 \leq k \leq 2n + 1$,

$$q \cdot \langle f, g \rangle = q \cdot \langle l_k(f), l_{k+1}(g) \rangle = \langle l_k(f), l_k(g) \rangle$$

Proof. The statement holds since l_k is linear and the conditions are linear in j and g . $\square$

Proposition 7.4. If $\alpha \in D_{n+1}$ and $\beta \in D_n$ then for $1 \leq k \leq 2n$, $q^c \langle \tau_k(\alpha), \beta \rangle_n = \langle (\alpha), l_k(\beta) \rangle_{n+1}$ where $c = 1$ if α has an inner loop from k to $k+1$ and $c = 0$ otherwise.

Proof. Consider the join $\alpha \cup l_k(\beta)y$. $l_k(\beta)$ has an inner loop from k to $k+1$ so the lower part of the gluing at the interval $[k, k+1]$ resembles the letter **U**, with its bottom half below the x-axis. Push the **U** onto the x-axis obtaining the figure $\sqcup$ with its base resting along the x-axis. The resulting chord diagram is none other than $\tau_k(\alpha) \cup \beta$ unless β originally had an arc from k to $k+1$. In that case there is an extra loop in the right expression. $\square$

8. CONSTRUCTION OF AN ORTHOGONAL BASIS FOR V_n

We now begin the construction our desired orthogonal basis elements using a two-level recursion. The outer recursion is on n itself, the size of the chord diagrams, the inner recursion is on the coordinate-wise partial ordering of restricted sequences associated to the chord diagrams. We now define our recursive expression for calculating the orthogonal basis elements (for $2 \leq a_n \leq n$):

$$(8.1) \quad e'_{(a_n, \dots, a_1)} = l_{a_n}(e'_{(a_{n-1}, \dots, a_1)}) - \frac{\Delta_{a_n-2}}{\Delta_{a_n-1}} \cdot e'_{(a_n-1, a_{n-1}, \dots, a_1)}$$

Note that this definition requires a recursion on n as well as the coordinate-wise partial order, $\preccurlyeq$, on D_n .

We initialize our recursion with the definition:

$$e'_{(1)} := e_{(1)}$$

With the convention $\Delta_{-1} = 0$, the recursive formula makes sense when $a_n = 1$ (although $(0, a_{n-1}, \dots, a_1)$ is an invalid restricted sequence) and simplifies to:

$$e'_{(1, a_{n-1}, \dots, a_1)} = l_1(e'_{(a_{n-1}, \dots, a_1)})$$

which is a recursion solely on n .

Remark 8.1. We can use the map l_1 to embed V_{n-1} into V_n (this sends the restricted sequence $(a_{n-1}, a_{n-2}, \dots, a_1)$ to $(1, a_{n-1}, a_{n-2}, \dots, a_1)$). The Markov form is scaled by q arising from the one additional component (by Prop 7.1).

9. QUAD MOVES

In this section, we investigate the relationship between the operation l_k and the partial ordering $\preccurlyeq$.

Lemma 9.1. Let α and $\beta \in D_n$. If $1 \leq k \leq 2n+1$ then:


$$(9.1) \quad \alpha \preccurlyeq \beta \Leftrightarrow l_k(\alpha) \preccurlyeq l_k(\beta)$$

The proof, given at the end of the section, will depend on an operation on chord diagrams called the *quad moves* — equivalently *θ -surgery* on a chord diagram.

Definition 9.1. For each pair of standard arcs, $\overline{ik}$ and $\overline{jl}$ in a chord diagram α with endpoints satisfying $i < k < j < l$ and such that no arc with endpoint in the interval $[k, j]$ separates the given arc, we define a quad move, $Quad_{i,j}(\alpha)$, to be the chord diagram obtained from α by removing the given pair and replacing them by the arcs $\overline{jk}$ and $\overline{il}$.

In the dual *rooted plane tree*, the condition on the arcs requires the corresponding edges to have a common parent node.

There is an important relationship between quad moves and the partial ordering. First, we demonstrate that the result of a quad move on a diagram is smaller in the partial order, $\preceq$, than the original diagram. Furthermore, we show for $\alpha \preceq \beta$ in D_n there is a sequence of quad moves which transforms β into α .



Definition 9.2. The *depth* of an arc a in $\alpha \in D_n$ is the number of arcs that lie over it. An arc a lies over another arc b if the left endpoint of a lies to the left of left endpoint b and the right endpoint of b lies to the right of the right endpoint of b .

Remark 9.1. The depth of the i th arc in $\alpha \in D_n$ is a_i where $(a_n, \dots, a_1)$ is the restricted sequence of α . Additionally, from the *rooted plane tree* perspective, depth is the distance to the root.

Proposition 9.1. If $(a_n, \dots, a_1) = \alpha \in D_n$ and $Q_{k,k'}$ a quad move of α involving arcs a_k and $a'_{k'}$ with $k < k'$, then $\beta = Q_{k,k'}(\alpha) \preceq \alpha$.

Proof. We show that the depth of the i th arc of $\beta = (b_n, \dots, b_1)$ is less than or equal to depth of the i th arc of α . The depth of the j th arc for $j < k'$ remains unchanged since they lie entirely to the right of the arcs involved, so $b_j = a_j$. Let $\{a_{k'+1}, \dots, a_{k''}\}$ be the arcs that lie under $a_{k'}$. Then clearly for $k' + 1 < j < k''$ we must have $a_{k'} < a_j$ and also that $b_{j-1} = a_j - 2$. But this implies that $b_{j-1} = a_j - 2 < a_j - 1 \leq a_{j-1}$. Thus, for $1 \leq j \leq k'' - 1$ we have $b_j \leq a_j$. Now the depth of the

k'' th arc of β is one less than the depth of the k' th arc of α , so $b_{k''} = a_{k'} - 1$. But then $b_{k''} = a_{k'} - 1 < a_{k'} < a_{k''}$. The depths of the other arcs remain unchanged under the quad move since they lie entirely to the left of it. $\square$

Proposition 9.2. Let α and β be in D_n . If $\alpha \preceq \beta$, then there is a sequence of quad moves which transforms β into α .

Proof. Note that removing an inner arc from a chord diagram does not affect the depth of the other arcs. In terms of restricted sequences, the equivalent way to express the deletion of an inner arc, which say is the i th arc in the diagram, would be to delete the i th component of the restricted sequence where the i th arc is an inner arc and shift all components to the left of the i th component to the right one space, thus obtaining an $n - 1$ restricted sequence. By the condition $\alpha \preceq \beta$, α 's rightmost inner arc lies to the right of β 's. Performing quad moves on β involving its rightmost inner arc and the arc immediately to its right, we may *pass the inner arc under* however many arcs on its right, until it aligns with the inner arc of β . This quad move leaves the depth of all arcs unaltered except for the depth of the right most inner arc which is lowered by one. The quad move leaves the ordering of the arcs unchanged except for interchanging the rightmost inner arc with the arc immediately to its left. Thus the quad move of β is still $\preceq \alpha$. So now we can assume that the rightmost inner arc of β is aligned with the rightmost inner arc of α . Let $\alpha', \beta' \in D_{n-1}$ be the chord diagrams obtained by removing the rightmost inner arcs from α and β respectively. Because the rightmost inner arcs are in the same position, we must have $\alpha' \preceq \beta'$. Then by induction on the size of chord diagrams there is a sequence of quad moves which take β' to α' . (The induction hypothesis is trivially satisfied for one chord diagrams.) Combining the two sequences of quad moves completes the proof. $\square$

Proof. To prove the initial lemma, we need only verify that quad moves commute with inserting inner arcs and quad moves also commute with deleting inner arcs, as long as the inner arc is not the result of a quad move. $\square$

We now give two corollaries of the relationship between the partial order and the l_k operation.

Corollary 9.1. The new basis $e'_{(a_n, \dots, a_1)}$ for the space, V_n is a linear combination of the original basis elements such that the coefficient of $e_{(b_n, \dots, b_1)}$ is nonzero in the linear combination iff $(b_n, \dots, b_1) \preceq (a_n, \dots, a_1)$.

Proof. $e'_{(a_n, \dots, a_1)} = l_{a_n}(e'_{(a_{n-1}, \dots, a_1)}) - \frac{\Delta_{a_n-2}}{\Delta_{a_n-1}} \cdot e'_{(a_{n-1}, a_{n-1}, \dots, a_1)}$. It suffices to show that $e'_{(a_{n-1}, a_{n-1}, \dots, a_1)}$ is a linear combination of basis elements in the *downset* of $\alpha = (a_n, \dots, a_1)$, the set $D(\alpha) := \{\beta \mid \beta \preceq \alpha\}$. Inductively $e'_{(a_{n-1}, a_{n-1}, \dots, a_1)}$ is a linear combination of basis elements $D(a_{n-1}, \dots, a_1)$. But by the transitivity of the partial ordering, this implies these basis elements are in $D(a_n, \dots, a_1)$. For the other term, we may inductively assume that $e'_{(a_{n-1}, \dots, a_1)}$ is a linear combination of elements all $D(a_{n-1}, \dots, a_1)$. Noticing that $l_{a_n}(e_{(a_{n-1}, \dots, a_1)}) = e_{(a_n, \dots, a_1)}$ we conclude from the preceding lemma, Lemma 9.1 that for any basis element e such that $e \preceq e_{(a_{n-1}, \dots, a_1)}$ we have $l_{a_n}(e) \preceq e_{(a_n, \dots, a_1)}$. The result follows. $\square$

Finally we determine the coefficient of the largest term (for $\preceq$) for the new basis.

Corollary 9.2. The change of basis is triangular, that is, in the expression of $e'_{(a_n, \dots, a_1)}$ as a linear combination of elements of the original basis, the polynomial coefficient of $e_{(a_n, \dots, a_1)}$ is one.

Proof. Note that only $l_{a_n}(e'_{(a_{n-1}, \dots, a_1)})$ has a nonzero $e_{(a_n, \dots, a_1)}$ component. Inductively we assume that $e'_{(a_{n-1}, \dots, a_1)}$ written as a linear combination of the original basis elements has 1 for the coefficient of $e_{(a_{n-1}, \dots, a_1)}$. By the conditions for equality in Lemma 9.1, the result follows. $\square$

10. ORTHOGONALITY

Theorem 10.1. The following two results hold:

- $(b_n, \dots, b_1) < (a_n, \dots, a_1) \Rightarrow \langle e_{(b_n, \dots, b_1)}, e'_{(a_n, \dots, a_1)} \rangle = 0$
- $(b_n, \dots, b_1) = (a_n, \dots, a_1) \Rightarrow \langle e_{(a_n, \dots, a_1)}, e'_{(a_n, \dots, a_1)} \rangle = \frac{\Delta_{a_n}}{\Delta_{a_n-1}} \dots \frac{\Delta_{a_1}}{\Delta_{a_1-1}}$

Proof. Our proof is by a double induction, an outer induction on the number of arcs, n , and an inner induction on the ordering of restricted sequences of length n .

We first note that the result holds for $n=1$, $\langle e_1, e'_1 \rangle = q = \frac{\Delta_1}{\Delta_{1-1}}$.

Our base case for the induction on the ordering is: Let $(b_n, \dots, b_1) \leq (a_n, \dots, a_1)$ with $a_n = 1$. Then we must have $b_n = 1$ and $(b_{n-1}, \dots, b_1) \leq (a_{n-1}, \dots, a_1)$ with equality iff we have equality in the original restricted n -sequences. Applying the recursion formula, we obtain:

$$\begin{aligned} \langle e_{(b_n, \dots, b_1)}, e'_{(a_n, \dots, a_1)} \rangle &= \langle e_{(1, b_{n-1}, \dots, b_1)}, e'_{(1, a_{n-1}, \dots, a_1)} \rangle \\ &= \langle e_{(1, b_{n-1}, \dots, b_1)}, l_1(e'_{(a_{n-1}, \dots, a_1)}) \rangle \\ &\quad - \frac{\Delta_{-1}}{\Delta_0} \langle e_{(1, b_{n-1}, \dots, b_1)}, e'_{(a_{n-1}, a_{n-1}, \dots, a_1)} \rangle \end{aligned}$$

the right term is zero since $\Delta_{-1} := 0$

$$\begin{aligned} &= q \cdot \langle \tau_1(e_{(1, b_{n-1}, \dots, b_1)}), e'_{(a_{n-1}, \dots, a_1)} \rangle \\ &= \frac{\Delta_1}{\Delta_0} \cdot \langle e_{(b_{n-1}, \dots, b_1)}, e'_{(a_{n-1}, \dots, a_1)} \rangle. \end{aligned}$$

By the adjoint relation between the operations and the Markov form, we have completed the proof in our base case.

The induction step is broken into four cases. While the calculations are different for each case, they are of similar flavor. For the sake of brevity, we give a complete calculation for a relatively complicated case, **Case 2(ii)** and sketch the others.

Case 1: $b_n < a_n - 1$:

$$\langle e_{(b_n, \dots, b_1)}, l_{a_n}(e'_{(a_{n-1}, \dots, a_1)}) \rangle = (q^c) \langle \tau_{a_n}(e_{(b_n, \dots, b_1)}), e'_{(a_{n-1}, \dots, a_1)} \rangle$$

which by the second part of Proposition 6.1

$$= (q^c) \langle (e_{(b'_{n-1}, \dots, b'_1)}), e'_{(a_{n-1}, \dots, a_1)} \rangle$$

for some restricted sequence $(b'_{n-1}, \dots, b'_1)$ with $b'_{n-1} \leq b_n$. And since $b'_{n-1} \leq b_n < a_n - 1 \leq a_{n-1}$ by induction on n ,

$$= (q^c) \langle e_{(b'_{n-1}, \dots, b'_1)}, e'_{(a_{n-1}, \dots, a_1)} \rangle = 0.$$

The monomial coefficient c is non-trivial in the case of an inner arc in $(b_n, \dots, b_1)$ at a_n .

For the rest of the cases, we use the following lemma:

Lemma 10.1. Let $(b_n, \dots, b_1), (a_{n-1}, \dots, a_1)$ be restricted sequences. If $j \leq b_n - 2 < a_{n-1}$ then $\langle e_{(b_n, \dots, b_1)}, e'_{(j, \dots, a_1)} \rangle = 0$.

Proof. Induction on j . For $j = 1$,

$$\begin{aligned} \langle e_{(b_n, \dots, b_1)}, e'_{(j, \dots, a_1)} \rangle &= \langle e_{(b_n, \dots, b_1)}, l_1(e'_{(a_{n-1}, \dots, a_1)}) \rangle \\ &= \langle \tau_1(e_{(b_n, \dots, b_1)}), e'_{(a_{n-1}, \dots, a_1)} \rangle \end{aligned}$$

By the first part of Proposition 6.1, $\tau_1(b_n, \dots, b_1) = (b'_{n-1}, \dots, b'_1)$ where $b'_{n-1} = b_n - 2 < a_{n-1}$. By the induction hypothesis on n ,

$$\begin{aligned} &= \langle \tau_1(e_{(b_n, \dots, b_1)}), e'_{(a_{n-1}, \dots, a_1)} \rangle \\ &= 0. \end{aligned}$$

For $1 < j \leq b_n - 2$,

$$\langle e_{(b_n, \dots, b_1)}, e'_{(j, \dots, a_1)} \rangle = \langle \tau_j(e_{(b_n, \dots, b_1)}), e'_{(a_{n-1}, \dots, a_1)} \rangle - \frac{\Delta_{j-2}}{\Delta_{j-1}} \cdot \langle e_{(b_n, \dots, b_1)}, e'_{(j-1, a_{n-1}, \dots, a_1)} \rangle$$

By induction on j , $\langle e_{(b_n, \dots, b_1)}, e'_{(j-1, a_{n-1}, \dots, a_1)} \rangle = 0$. Once again, finding that $\tau_j(e_{(b_n, \dots, b_1)}) = (b'_{n-1}, \dots, b'_1)$ with $b'_{n-1} \leq b_n - 2$ we conclude by the induction hypothesis on n that the first term is also zero.

□

Case 2: $b_n = a_n - 1$

$$\begin{aligned} \langle e_{(b_n, \dots, b_1)}, e'_{(a_n, \dots, a_1)} \rangle &= \langle e_{(b_n, \dots, b_1)}, l_{a_n}(e'_{(a_{n-1}, \dots, a_1)}) \rangle \\ &= \frac{\Delta_{a_n-2}}{\Delta_{a_n-1}} \langle e_{(b_n, \dots, b_1)}, e'_{(a_n-1, a_{n-1}, \dots, a_1)} \rangle \\ &= \langle e_{(b_{n-1}, \dots, b_1)}, e'_{(a_{n-1}, \dots, a_1)} \rangle \\ &= \frac{\Delta_{a_n-2}}{\Delta_{a_n-1}} \langle e_{(b_n, \dots, b_1)}, e'_{(a_n-1, a_{n-1}, \dots, a_1)} \rangle \end{aligned}$$

Case 2(i): $(b_n, \dots, b_1) = (a_n - 1, a_{n-1}, \dots, a_1)$ By the induction hypothesis we have:

$$\begin{aligned} &= \left(\frac{\Delta_{a_n-1}}{\Delta_{a_n-1-1}} \cdot \frac{\Delta_{a_n-2}}{\Delta_{a_n-2-1}} \cdots \frac{\Delta_{a_1}}{\Delta_{a_1-1}} \right) - \frac{\Delta_{a_n-2}}{\Delta_{a_n-1}} \cdot \left(\frac{\Delta_{a_n-1}}{\Delta_{a_n-2}} \cdot \frac{\Delta_{a_n-1}}{\Delta_{a_n-1-1}} \cdot \frac{\Delta_{a_n-2}}{\Delta_{a_n-2-1}} \cdots \frac{\Delta_{a_1}}{\Delta_{a_1-1}} \right) \\ &= 0. \end{aligned}$$

Case 2(ii): $b_n = a_n - 1$ and $(b_{n-1}, \dots, b_1) \neq (a_{n-1}, \dots, a_1)$

Expanding $e'_{(a_n, \dots, a_1)}$ three times with the recursive definition, we obtain:

$$\begin{aligned} \langle e_{(b_n, \dots, b_1)}, e'_{(a_n, \dots, a_1)} \rangle &= \langle e_{(b_n, \dots, b_1)}, l_{a_n}(e'_{(a_{n-1}, \dots, a_1)}) \rangle \\ &\quad - \frac{\Delta_{a_n-2}}{\Delta_{a_n-1}} \langle e_{(b_n, \dots, b_1)}, l_{a_n-1}(e'_{(a_{n-1}, \dots, a_1)}) \rangle \\ &\quad + \frac{\Delta_{a_n-3}}{\Delta_{a_n-1}} \langle e_{(b_n, \dots, b_1)}, l_{a_n-2}(e'_{(a_{n-1}, \dots, a_1)}) \rangle \\ &\quad - \frac{\Delta_{a_n-4}}{\Delta_{a_n-1}} \langle e_{(b_n, \dots, b_1)}, e'_{(a_n-3, a_{n-1}, \dots, a_1)} \rangle \end{aligned}$$

and by the adjoint property of the operations in Proposition 7.3 we get

$$\begin{aligned} &= \langle \tau_{a_n}(e_{(b_n, \dots, b_1)}), e'_{(a_{n-1}, \dots, a_1)} \rangle \\ &\quad - q \cdot \frac{\Delta_{a_n-2}}{\Delta_{a_n-1}} \langle \tau_{a_n-1}(e_{(b_n, \dots, b_1)}), e'_{(a_{n-1}, \dots, a_1)} \rangle \\ &\quad + \frac{\Delta_{a_n-3}}{\Delta_{a_n-1}} \langle \tau_{a_n-2}(e_{(b_n, \dots, b_1)}), e'_{(a_{n-1}, \dots, a_1)} \rangle \\ &\quad - \frac{\Delta_{a_n-4}}{\Delta_{a_n-1}} \langle e_{(b_n, \dots, b_1)}, e'_{(a_n-3, a_{n-1}, \dots, a_1)} \rangle \end{aligned}$$

since $a_n - 2, a_n - 1, a_n$ equal $b_n - 1, b_n, b_n + 1$ respectively, we use Remark 5.1 to get

$$\begin{aligned} &= \left[\left(1 - q \frac{\Delta_{a_n-2}}{\Delta_{a_n-1}} + \frac{\Delta_{a_n-3}}{\Delta_{a_n-1}} \right) \cdot \langle e_{(b_{n-1}, \dots, b_1)}, e'_{(a_{n-1}, \dots, a_1)} \rangle \right] \\ &\quad - \left[\frac{\Delta_{a_n-4}}{\Delta_{a_n-1}} \cdot \langle e_{(b_n, \dots, b_1)}, e'_{(a_n-3, a_{n-1}, \dots, a_1)} \rangle \right]. \end{aligned}$$

By the recursion property for the Chebyshev polynomials 4.1, $(1 - q \frac{\Delta_{a_n-2}}{\Delta_{a_n-1}} + \frac{\Delta_{a_n-3}}{\Delta_{a_n-1}}) = 1 - 1 = 0$

For $j = a_n - 3$ we have $j = a_n - 3 = b_n + 1 - 3 = b_n - 2 < b_n = a_n - 1 \leq a_{n-1}$ hence the required conditions hold for applying Lemma 10.1 and we conclude that $\langle e_{(b_n, \dots, b_1)}, e'_{(a_n-3=j, a_{n-1}, \dots, a_1)} \rangle = 0$.

Some extra care should be taken when a_n is small. Because of the hypothesis, $b_n = a_n - 1$, we may assume $2 \leq a_n$. When $4 \leq a_n$ expanding by recursion is well defined. For $3 = a_n$ the expansion also holds, but there is no second term. The case $2 = a_n$ deserves a special treatment. Given $(1, b_{n-1}, \dots, b_1)$:

$$\begin{aligned}
\langle e_{(1,\dots,b_1)}, e'_{(2,\dots,a_1)} \rangle &= \langle l_1(e_{(b_{n-1},\dots,b_1)}), l_2(e'_{(a_{n-1},\dots,a_1)}) \rangle \\
&\quad - \frac{\Delta_0}{\Delta_1} \langle e_{(1,b_{n-1},\dots,b_1)}, e'_{(1,a_{n-1},\dots,a_1)} \rangle \\
&= \langle l_1(e_{(b_{n-1},\dots,b_1)}), l_2(e'_{(a_{n-1},\dots,a_1)}) \rangle \\
&\quad - \frac{\Delta_0}{\Delta_1} \langle l_1(e_{(b_{n-1},\dots,b_1)}), l_1(e'_{(a_{n-1},\dots,a_1)}) \rangle
\end{aligned}$$

And by Proposition 7.3 we get

$$\begin{aligned}
&= \langle e_{(b_{n-1},\dots,b_1)}, e'_{(a_{n-1},\dots,a_1)} \rangle \\
&\quad - \frac{1}{q} \langle e_{(b_{n-1},\dots,b_1)}, e'_{(a_{n-1},\dots,a_1)} \rangle \\
&= 0.
\end{aligned}$$

Case 3: $b_n = a_n$ and $(b_{n-1}, \dots, b_1) < (a_{n-1}, \dots, a_1)$

When $1 = a_n$ the following holds by Proposition 7.3 and by induction on n :

$$\begin{aligned}
\langle e_{(b_n,\dots,b_1)}, e'_{(a_n,\dots,a_1)} \rangle &= \langle l_1(e_{(b_{n-1},\dots,b_1)}), l_1(e'_{(a_{n-1},\dots,a_1)}) \rangle \\
&= q \cdot \langle e_{(b_{n-1},\dots,b_1)}, e'_{(a_{n-1},\dots,a_1)} \rangle \\
&= 0
\end{aligned}$$

For $2 = a_n$ we expand twice and apply induction on n .

$$\begin{aligned}
\langle e_{(b_n,\dots,b_1)}, e'_{(a_n,\dots,a_1)} \rangle &= \langle l_2(e_{(b_{n-1},\dots,b_1)}), l_2(e'_{(a_{n-1},\dots,a_1)}) \rangle \\
&\quad - \frac{\Delta_0}{\Delta_1} \langle l_2(e_{(b_{n-1},\dots,b_1)}), l_1(e'_{(a_{n-1},\dots,a_1)}) \rangle \\
&= q \cdot \langle e_{(b_{n-1},\dots,b_1)}, e'_{(a_{n-1},\dots,a_1)} \rangle - \frac{1}{q} \langle e_{(b_{n-1},\dots,b_1)}, e'_{(a_{n-1},\dots,a_1)} \rangle \\
&= 0
\end{aligned}$$

Lastly for $2 \leq a_n$ we expand twice obtaining an expression similar to the one above:

$$\begin{aligned}
&= \langle l_{b_n=a_n}(e_{(b_{n-1},\dots,b_1)}), l_{a_n}(e'_{(a_{n-1},\dots,a_1)}) \rangle \\
&\quad - \frac{\Delta_{a_n-2}}{\Delta_{a_n-1}} \langle l_{b_n=a_n}(e_{(b_{n-1},\dots,b_1)}), l_{a_n-1}(e'_{(a_{n-1},\dots,a_1)}) \rangle \\
&\quad + \frac{\Delta_{a_n-3}}{\Delta_{a_n-1}} \langle e_{(b_n,\dots,b_1)}, e'_{(a_n-2,a_{n-1},\dots,a_1)} \rangle \\
&= (q - \frac{\Delta_{a_n-2}}{\Delta_{a_n-1}}) \cdot \langle e_{(b_{n-1},\dots,b_1)}, e'_{(a_{n-1},\dots,a_1)} \rangle \\
&\quad + \frac{\Delta_{a_n-3}}{\Delta_{a_n-1}} \cdot \langle e_{(b_n,\dots,b_1)}, e'_{(a_n-2,a_{n-1},\dots,a_1)} \rangle
\end{aligned}$$

Note, by induction on n , that $\langle e_{(b_{n-1}, \dots, b_1)}, e'_{(a_{n-1}, \dots, a_1)} \rangle = 0$

$$= \frac{\Delta_{a_n-3}}{\Delta_{a_n-1}} \cdot \langle e_{(b_n, \dots, b_1)}, e'_{(a_n-2, a_{n-1}, \dots, a_1)} \rangle$$

For $j = a_n - 2$ we have $j = a_n - 2 = b_n - 2 < b_n - 1 = a_n - 1 \leq a_{n-1}$ hence we can conclude from Lemma 10.1 that

$$\begin{aligned} &= \langle e_{(b_n, \dots, b_1)}, e'_{(a_n-2, a_{n-1}, \dots, a_1)} \rangle \\ &= 0 \end{aligned}$$

Case 4: $(b_n, \dots, b_1) = (a_n, \dots, a_1)$

When $a_n = b_n = 1$ the base case gives the desired conclusion. If $a_n = 2$ we expand twice by recursion to obtain:

$$\begin{aligned} \langle e_{(b_n, \dots, b_1)}, e'_{(a_n, \dots, a_1)} \rangle &= \langle l_2(e_{(b_{n-1}, \dots, b_1)}), l_2(e'_{(a_{n-1}, \dots, a_1)}) \rangle \\ &\quad - \frac{\Delta_0}{\Delta_1} \cdot \langle l_2(e_{(b_{n-1}, \dots, b_1)}), l_1(e'_{(a_{n-1}, \dots, a_1)}) \rangle \\ &= (q - \frac{\Delta_0}{\Delta_1}) \cdot \langle e_{(b_{n-1}, \dots, b_1)}, e'_{(a_{n-1}, \dots, a_1)} \rangle \\ &= \frac{\Delta_2}{\Delta_1} \cdot \frac{\Delta_{a_n-1}}{\Delta_{a_n-1}-1} \dots \frac{\Delta_{a_1}}{\Delta_{a_1}-1} \end{aligned}$$

which is the desired expression. For $3 \leq a_n$ we again expand twice, but an extra term results.

$$\begin{aligned} &= \langle l_{a_n=b_n}(e_{(b_{n-1}, \dots, b_1)}), l_{a_n=b_n}(e'_{(a_{n-1}, \dots, a_1)}) \rangle \\ &\quad - \frac{\Delta_{a_n-2}}{\Delta_{a_n-1}} \cdot \langle l_{b_n=a_n}(e_{(b_{n-1}, \dots, b_1)}), l_{a_n-1}(e'_{(a_{n-1}, \dots, a_1)}) \rangle \\ &\quad + \frac{\Delta_{a_n-3}}{\Delta_{a_n-1}} \langle e_{(b_n, \dots, b_1)}, e'_{(a_n-2, \dots, a_1)} \rangle \end{aligned}$$

Setting $j = a_n - 2$ we have $j = a_n - 2 = b_n - 2 < b_n - 1 = a_n - 1 \leq a_{n-1}$ we may apply Lemma 10.1 to obtain:

$$= \langle e_{(b_n, \dots, b_1)}, e'_{(a_n-2, \dots, a_1)} \rangle = 0$$

The remaining terms simplify to

$$\begin{aligned} &= \langle l_{a_n=b_n}(e_{(b_{n-1}, \dots, b_1)}), l_{a_n=b_n}(e'_{(a_{n-1}, \dots, a_1)}) \rangle \\ &\quad - \frac{\Delta_{a_n-2}}{\Delta_{a_n-1}} \cdot \langle l_{b_n=a_n}(e_{(b_{n-1}, \dots, b_1)}), l_{a_n-1}(e'_{(a_{n-1}, \dots, a_1)}) \rangle \\ &= (q - \frac{\Delta_{a_n-2}}{\Delta_{a_n-1}}) \langle e_{(b_{n-1}, \dots, b_1)}, e'_{(a_{n-1}, \dots, a_1)} \rangle \\ &= \frac{\Delta_{a_n}}{\Delta_{a_n-1}} \langle e_{(b_{n-1}, \dots, b_1)}, e'_{(a_{n-1}, \dots, a_1)} \rangle \end{aligned}$$

and by induction on n we obtain what we want

$$= \frac{\Delta_{a_n}}{\Delta_{a_n-1}} \cdot \frac{\Delta_{a_{n-1}}}{\Delta_{a_{n-1}-1}} \cdots \frac{\Delta_{a_1}}{\Delta_{a_1-1}}$$

□

11. THE MARKOV FORM IN THE ORTHOGONAL BASIS

We now prove our main result that the space of chord diagrams, V_n , equipped with the Markov symmetric bilinear form, $\langle \cdot, \cdot \rangle_n$, has $\{e'_\alpha\}$ for an orthogonal basis.

Theorem 11.1. The basis $\{e'_\alpha\}$ is orthogonal. Furthermore, for $\alpha = (a_n, \dots, a_1)$.

- $\langle e'_\alpha, e'_\alpha \rangle = \frac{\Delta_{a_n}}{\Delta_{a_n-1}} \cdots \frac{\Delta_{a_1}}{\Delta_{a_1-1}}$
- If $\beta \neq \alpha$ then $\langle e'_\beta, e'_\alpha \rangle = 0$

Proof. By symmetry, we may assume $\beta = (b_n, \dots, b_1) \leq \alpha = (a_n, \dots, a_1)$. By the corollary, we can write $e'_{(b_n, \dots, b_1)} = \sum_{\alpha \preceq \in D_n} P_\alpha(q) \alpha$ where the $P_\alpha(q)$ are rational polynomials which are 0 for $\alpha \not\leq (b_n, \dots, b_1)$.

Hence the computation of the Markov form yields: $\langle e'_{(b_n, \dots, b_1)}, e'_{(a_n, \dots, a_1)} \rangle = \sum_{\alpha \preceq (b_n, \dots, b_1) \in D_n} P_\alpha(q) \langle \alpha, e'_{(a_n, \dots, a_1)} \rangle$.

In the case $(b_n, \dots, b_1) < (a_n, \dots, a_1)$, we have for every α in the sum: $\alpha \preceq (b_n, \dots, b_1) \Rightarrow \alpha \leq (b_n, \dots, b_1) \Rightarrow \alpha < (a_n, \dots, a_1)$. So by Theorem 10.1, each bracket term is 0, so the sum is 0.

When $(b_n, \dots, b_1) = (a_n, \dots, a_1)$ the same reasoning applies, except that there is a single term left, the one term where equality holds:

$$P_{\alpha=(a_n, \dots, a_1)}(q) \langle e_{(a_n, \dots, a_1)}, e'_{(a_n, \dots, a_1)} \rangle$$

But, as previously remarked, $P_{(a_n, \dots, a_1)}(q) = 1$ and by 10.1 only the desired product $\frac{\Delta_{a_n}}{\Delta_{a_n-1}} \cdots \frac{\Delta_{a_1}}{\Delta_{a_1-1}}$ remains. □

12. RELATION WITH OTHER ORTHOGONAL BASES

The following technique for constructing orthogonal bases for quantum modules, implicit in the literature, was shown to the authors by Pat Gilmer. The method uses the calculus of trivalent graphs for computing three-manifold quantum invariants (see Kauffman-Lins[5]).

The method begins with a fixed choice of a trivalent plane tree in the unit disk with $2n$ terminal (external) edges and $2n - 3$ internal edges. In the method of Kauffman and Lins, the edges must be *coloured* with (quantum) integers satisfying all the triangular inequalities (at each trivalent vertex the sum of the integers must be even and the sum of any two integers minus the third must be non-negative). For this construction all the external edges must be coloured one. The number of such colorings is the Catalan number, as expected.

Next, we pair two such colorations of our chosen trivalent plane tree by gluing the univalent ends and computing the resulting quantum invariant in $\mathbb{Z}[q^\epsilon]$. Using two-spheres meeting the paired configuration in two points labelled one, we inductively compute the Markov pairing using the fusion rule (Lemma 7, p. 40[5]). The components of the sum either all vanish (when the colorations are different) or one

single summand remains, which is a product of theta coefficients, easily rewritten as a quotient of Chebyshev polynomials.

For $n = 3$, there are three non-isomorphic configurations (removing the six terminal edges from a plane trivalent tree on 9 edges leaves a tree with three edges which must either be a linear chain or a trivalent **Y**). The **Y** configuration completes to a unique plane trivalent tree with 9 edges and the chain completes to two plane trivalent trees, differing in whether or not, the terminal edges connected to the interior vertices of the chain are on the same or opposite *sides*). The associated change of basis to the *non-crossing* chord diagram basis (after making a choice of root for the chord diagram) and the corresponding orthogonal matrices are given below.

12.1. Colored Trivalent Graphs. First, for the **Y** configuration, we provide the change of basis matrix:

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ -\frac{1}{q} & 0 & 1 & 0 & 0 \\ -\frac{1}{q} & 0 & 0 & 0 & 1 \\ -\frac{1}{q} & 1 & 0 & 0 & 0 \\ \frac{2}{q^2} & -\frac{1}{q} & -\frac{1}{q} & 1 & -\frac{1}{q} \end{pmatrix}$$

and the Markov form in this orthogonal basis.

$$\begin{pmatrix} q^3 & 0 & 0 & 0 & 0 \\ 0 & (q-1)q(q+1) & 0 & 0 & 0 \\ 0 & 0 & (q-1)q(q+1) & 0 & 0 \\ 0 & 0 & 0 & (q-1)q(q+1) & 0 \\ 0 & 0 & 0 & 0 & \frac{(q-1)(q+1)(q^2-2)}{q} \end{pmatrix}$$

Next the same information is provided for the *same side* configuration:

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ -\frac{1}{q} & 1 & 0 & 0 & 0 \\ -\frac{1}{q} & 0 & 1 & 0 & 0 \\ \frac{1}{q^2} & -\frac{1}{q} & -\frac{1}{q} & 1 & 0 \\ -\frac{q}{q^2-1} & \frac{1}{q^2-1} & \frac{1}{q^2-1} & -\frac{q}{q^2-1} & 1 \end{pmatrix}$$

$$\begin{pmatrix} q^3 & 0 & 0 & 0 & 0 \\ 0 & (q-1)q(q+1) & 0 & 0 & 0 \\ 0 & 0 & (q-1)q(q+1) & 0 & 0 \\ 0 & 0 & 0 & \frac{(q-1)^2(q+1)^2}{q} & 0 \\ 0 & 0 & 0 & 0 & q(q^2-2) \end{pmatrix}$$

and, finally for the opposite side configuration:

$$\begin{pmatrix} 0 & 0 & 1 & 0 & 0 \\ 1 & 0 & -\frac{1}{q} & 0 & 0 \\ 0 & 0 & -\frac{1}{q} & 1 & 0 \\ -\frac{1}{q} & 1 & q^{-2} & -\frac{1}{q} & 0 \\ -\frac{q}{-1+q^2} & \frac{1}{-1+q^2} & \frac{1}{-1+q^2} & \frac{q}{-1+q^2} & 1 \end{pmatrix}$$

$$\begin{pmatrix} q^3 & 0 & 0 & 0 & 0 \\ 0 & (q-1)q(q+1) & 0 & 0 & 0 \\ 0 & 0 & (q-1)q(q+1) & 0 & 0 \\ 0 & 0 & 0 & \frac{(q-1)^2(q+1)^2}{q} & 0 \\ 0 & 0 & 0 & 0 & -2q+q^3 \end{pmatrix}$$

12.2. Computation for the poset recursion construction. Here is the change of basis matrix for the poset-recursion.

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ -\frac{1}{q} & 1 & 0 & 0 & 0 \\ -\frac{1}{q^2} & 0 & 1 & 0 & 0 \\ \frac{1}{q^2} & -\frac{1}{q} & -\frac{1}{q} & 1 & 0 \\ -\frac{q}{q^2-1} & \frac{1}{q^2-1} & \frac{1}{q^2-1} & -\frac{q}{q^2-1} & 1 \end{pmatrix}$$

and the matrix for the Markov form.

$$\begin{pmatrix} q^3 & 0 & 0 & 0 & 0 \\ 0 & (q-1)q(q+1) & 0 & 0 & 0 \\ 0 & 0 & (q-1)q(q+1) & 0 & 0 \\ 0 & 0 & 0 & \frac{(q-1)^2(q+1)^2}{q} & 0 \\ 0 & 0 & 0 & 0 & q(q^2-2) \end{pmatrix}$$

The *same-side* configuration gives the same diagonal form and triangular change of basis as the recursively computed diagonalized matrix. Although the orthogonal basis is the same, the poset-recursion formula provides an explanation for the location of the non-zero coefficients in the triangular change of basis: the non-zero coefficients can only occur in positions corresponding to those restricted sequences smaller than the given one, that is, in the down-set of the given element in the coordinate-wise partial order.

13. FURTHER DEVELOPMENT

In a subsequent paper, we will develop further properties of this poset from the perspective of *non-crossing fixed point free involutions* in a permutation group of a set of even cardinality. In addition, this new viewpoint will allow us to extend this theory to annular braids, that is, Artin braids of type $\mathbb{B}_n$.

REFERENCES

- [1] Blanchet, C. and Habegger, N. and Masbaum, G. and Vogel, P., *Three-manifold invariants derived from the Kauffman bracket*, Topology, Vol. 31, No. 4, 1992, p. 685–699.
- [2] R. A. Brualdi, *Introductory Combinatorics*, North-Holland, New York, 1977.
- [3] Patrick M. Gilmer and John M. Harris, *On the Kauffman bracket skein module of the quaternionic manifold*, *arXiv:math.GT/0406152*
- [4] Jones, V. F. R., *Index for subfactors*, Inventiones Mathematicae, Vol. 72, No. 1, 1983, p. 1–25.
- [5] Kauffman, Louis H. and Lins, Sóstenes L., *Temperley-Lieb recoupling theory and invariants of 3-manifolds*, Annals of Mathematics Studies, vol. 134, Princeton University Press, 1994, ISBN0-691-03640-3.
- [6] Ki Hyoung Ko and Lawrence Smolinsky, *A Combinatorial Matrix in 3-Manifold Theory*, Pacific Journal of Mathematics Vol. 149, No. 2, 1991, 319–336.

- [7] W.B.R. Lickorish, *3-Manifold Invariants from the Combinatorics of the Jones Polynomial*, Pacific Journal of Mathematics Vol. 149, No. 2, 1991, 337-347.
- [8] N. Y. Reshetikhin and V. G. Turaev, *Invariants of 3-Manifolds Via Link Polynomials and Quantum Groups* Invent. Math., 103, 1991, 547-597.
- [9] Stanley, Richard P., *Enumerative combinatorics. Vol. 2*, Cambridge University Press, 1999, 581 p., ISBN: 0-521-56069-1
- [10] E. Witten, *Quantum Field Theory and Jones' Polynomial*, Comm. Math. Phys., 121, 1989, 351-399.

DEPT. OF MATH., STANFORD UNIVERSITY, STANFORD, CA
E-mail address: `genauer@math.stanford.edu`

DEPARTMENT OF MATHEMATICS, LOUISIANA STATE UNIVERSITY, BATON ROUGE, LA 70803-4918
E-mail address: `stoltz@lsu.edu`